Neuroscience Letters, 117 (1990) 364–368
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abe, K., Xie, F., Watanabe, Y. and Saito, H., Glycine facilitates induction of long-term potentiation of evoked potential in rat hippocampus, 87
Ackerman, J.M. and White, F.J., A10 somatodendritic dopamine autoreceptor sensitivity following withdrawal from repeated cocaine treatment, 181
Amaducci, L., see Sorbi, S., 165
Amann, R., see Donnerer, J., 331
Antal, M., see Shupliakov, O.V., 289
Aoki, M., see Arai, T., 353
Arai, T., Aoki, M., Murakawa, M., Nakao, S., Mori, Kenjiro and Kurihara, N., The effects of halothane on the contents of putative transmitter amino acids in whole rat brain, 353
Armbrecht, H.J., see Strong, R., 175
Avoli, M., Epileptiform discharges and a synchronous GABAergic potential induced by 4-aminopyridine in the rat immature hippocampus, 93

Bansinath, M., Ramabadran, K., Turndorf, H. and Puig, M.M., Effects of the benzomorphan κ-opiate, MR 2266 and its (+) enantiomer MR 2267, on thermonociceptive reactions in different strains of mice, 212
Barraclough, C., see Rogers, A.T., 194
Berardelli, A., see Cruccu, G., 68
Berger, W., see Dietz, V., 81
Betschart, J., see Piot, O., 140
Blanchard, J.C., see Piot, O., 140
Bloch, B., see Moine, C.L., 248
Block, F., see Jaspers, R.M.A., 149
Brock, J.W., Ng, J.P. and Justice, J.B.Jr., Effect of chronic cocaine on dopamine synthesis in the nucleus accumbens as determined by microdialysis perfusion with NSD-1015, 234
Brodie, C. and Sampson, S.R., Thyroid hormones up-regulate Ca-channels in cultured skeletal muscle of the rat, 325
Bublitz, C., see Lasher, R.S., 43

Carboni, S., Melis, F., Pani, L., Hadjiconstantinou, M. and Rossetti, Z.L., The non-competitive NMDA-receptor antagonist MK-801 prevents the massive release of glutamate and aspartate from rat striatum induced by 1-methyl-4-phenylpyridinium (MPP), 129
Carpenter, D.O., see Evans, D.B.M.L., 117
Case, C.P., see Rogers, A.T., 194
Cespuglio, R., see Houdouin, F., 218
Chen, H., see Zidichouski, J.A., 123
Costentin, J., see Leroux-Nicollet, I., 1
Cowie, R.J., see McKenzie, J.C., 253
Cron, K.C., see Gadient, R.A., 335
Cruccu, G., Berardelli, A., Inghilleri, M. and Manfredi, M., Corticobulbar projections to upper and lower facial motoneurons. A study by magnetic transcranial stimulation in man, 68
Csillik, B., see Hajós, F., 8

Darchen, F., see Leroux-Nicollet, I., 1
Demontis, M.G., Fadda, P., Devoto, P., Martellotta, M.C. and Fratta, W., Sleep deprivation increases dopamine D_1 receptor antagonist [^{3}H]SCH 23390 binding and dopamine-stimulated adenylate cyclase in the rat limbic system, 224
Der Kloot, W.V., see Yu, S.P., 111
Devoto, P., see Demontis, M.G., 224
Dietz, V., Feuerstein, T.J. and Berger, W., Significance of dopamine receptor antagonists in human postural control, 81
Distefano, L.M., see Kish, S.J., 347
Donnerer, J. and Amann, R., Capsaicin-evoked neuropeptide release is not dependent on membrane potential changes, 331
Dozič, S., see Kish, S.J., 347

Erdélyi, L., Ilyin, V.I., Lozovaya, N.A. and Vulfius, C.A., Intracellular Mg^2 modulates the A-current and its blockage by catechol in isolated *Lymnaea* neurons, 99
Evans, D.B.M.L., Rahmann, H. and Carpenter, D.O., Zn^{2+} blocks the voltage activated calcium current of *Aplysia* neurons, 117

Facci, L., see Skaper, S.D., 154
Fadda, P., see Demontis, M.G., 224
Feuerstein, T.J., see Dietz, V., 81
Fischer, S., see Wolf, G., 207
Fratta, W., see Demontis, M.G., 224
Fujino, Yu., see Ono, T., 259

Gadient, R.A., Cron, K.C. and Otten, U., Interleukin-1 β and tumor necrosis factor-α synergis-

0304-3940/90/$ 03.50 © 1990 Elsevier Scientific Publishers Ireland Ltd.

tically stimulate nerve growth factor (NGF) release from cultured rat astrocytes, 335
Gähwiler, B.H., see Knöpfel, T., 295
Garcia, L., see Verdière-Sahuque, M., 341
Geiger, J.D., see Wan, W., 160
Geula, C., see Schätz, C.R., 56
Gharib, A., see Houdouin, F., 218
Goto, K., Kurihara, T., Takahashi, Y. and Kondo, H., Expression of genes for the myelin-specific proteins in oligodendrocytes in vivo demands the presence of axons, 269
Goudou, D., see Verdière-Sahuque, M., 341
Grafe, P., see Knöpfel, T., 295

Hadjiconstantinou, M., see Carboni, S., 129
Haga, S., see Kametani, F., 62
Hajós, F., Csillik, B. and Knyihár-Csillik, E., Alterations in glial fibrillary acidic protein immunoreactivity in the upper dorsal horn of the rat spinal cord in the course of transganglionic degenerative atrophy and regenerative proliferation, 8
Hale, C., see Strong, R., 175
Harrison, R., see Rogers, A.T., 194
Hartle, D.K., see Phelix, C.F., 31
Harvey, A.R., see Tan, M.M.L., 14
Hass, P., see Wolf, G., 207
Hećimović, H., see Randić, M., 74
Heim, C., see Jaspers, R.M.A., 149
Hernandez, L., see Rada, P., 144
H.N.Deck, J., see Kish, S.J., 347
Hofmann, H.-D., see Huba, R., 37
Hornykiewicz, O., see Kish, S.J., 347
Hösli, E., see Hösli, L., 307
Hösli, L., Hösli, E., Redle, S., Rojas, J. and Schramek, H., Action of baclofen, GABA and antagonists on the membrane potential of cultured astrocytes of rat spinal cord, 307
Houdouin, F., Cespuglio, R., Gharib, A., Sarda, N. and Jouvet, M., Time-course variations induced by pargyline on the 5-hydroxyindole compounds measured in the nucleus raphe dorsalis and in blood: a voltammetric and HPLC approach in the rat, 218
Hoyer, S., Nitsch, R. and Oesterreich, K., Ammonia is endogenously generated in the brain in the presence of presumed and verified dementia of Alzheimer type, 358
Huba, R. and Hofmann, H.-D., Identification of GABAergic amacrine cell-like neurons developing in chick retinal monolayer cultures, 37
Hulsebosch, C.E., see Urschel, B.A., 50
Hutchison, W.D., see Morton, C.R., 319

Ilyin, V.I., see Erdélyi, L., 99
Inagami, T., see McKenzie, J.C., 253
Inghilleri, M., see Cruccu, G., 68
Ishii, T., see Kametani, F., 62

Jaspers, R.M.A., Block, F., Heim, C. and Sontag, K.-H., Spatial learning is affected by transient occlusion of common carotid arteries (2VO): comparison of behavioural and histopathological changes after '2VO' and 'four-vessel-occlusion' in rats, 149
Jouvet, M., see Houdouin, F., 218
Justice, J.B.Jr., see Brock, J.W., 234

Kakei, S. and Shinoda, Y., Parietal projection of thalamocortical fibers from the ventroanterior-ventrolateral complex of the cat thalamus, 280
Kametani, F., Tanaka, K., Sato, M., Haga, S., Saitoh, T., Shinoda, T. and Ishii, T., A monoclonal antibody Hy20-54-16-3L to lambda (λ) light chain of human immunoglobulin reacts with amyloid in Alzheimer's disease brain, 62
Kameyama, T., see Nabeshima, T., 200
Kaneko, S., Sugimura, M. and Satoh, M., Interspecies differences in the population of multiple glutamate receptors expressed in *Xenopus* oocytes after injection of guinea pig, mouse and rat forebrain mRNAs, 134
Keilhoff, G., see Wolf, G., 207
Kish, S.J., Distefano, L.M., Dozič, S., Robitaille, Y., Rajput, Ali, J.H.N.Deck, and Hornykiewicz, O., [^{3}H]Vesamicol binding in human brain cholinergic deficiency disorders, 347
Kitao, Ya., see Kudo, M., 26
Kitazawa, S., Application of principal component analysis for characterizing convergence patterns of inputs in interneurones of the cat forelimb segments, 313
Knöpfel, T., Spuler, A., Grafe, P. and Gähwiler, B.H., Cytosolic calcium during glucose deprivation in hippocampal pyramidal cells of rats, 295
Knyihár-Csillik, E., see Hajós, F., 8
Kondo, H., see Goto, K., 269
Kondo, H., see Yamamoto, M., 275
Kudo, M., Kitao, Y. and Nakamura, Y., Differential organization of crossed and uncrossed projections from the superior olive to the inferior colliculus in the mole, 26
Kurata, K. and Shibata, R., Biphasic effect of locally applied apomorphine and 2,3,4,5-tetrahydro-7,8-dihydroxy-1-phenyl-1*H*-3-benzazepine on the release of striatal dopamine investigated by means of brain dialysis, 264

Kurihara, N., see Arai, T., 353
Kurihara, T., see Goto, K., 269

Lasher, R.S., Lutz, E.M., Mulholland, F., Sanderson, R., Stewart, J.M. and Bublitz, C., Immunocytochemical localization of endopeptidase-24.11 in the nucleus tractus solitarius of the rat brain, 43
Leon, A., see Skaper, S.D., 154
Leroux-Nicollet, I., Darchen, F., Scherman, D. and Costentin, J., Postnatal development of the monoamine vesicular transporter in mesencephalic and telencephalic regions of the rat brain: a quantitative autoradiographic study with [^{3}H]dihydrotetrabenazine, 1
Lipton, P., see Lobner, D., 169
Li, Yu.-Q., Rao, Z.-R. and Shi, J.-W., Collateral projections from the midbrain periaqueductal gray to the nucleus raphe magnus and nucleus accumbens in the rat. A fluorescent retrograde double-labelling study, 285
Lobner, D. and Lipton, P., σ-Ligands and non-competitive NMDA antagonists inhibit glutamate release during cerebral ischemia, 169
Lozovaya, N.A., see Erdélyi, L., 99
Lozsádi, D., see Réthelyi, M., 20
Lund, D.D., see Pardini, B.J., 300
Lunt, G.G., see Rogers, A.T., 194
Lutz, E.M., see Lasher, R.S., 43

Mailleux, P. and Vanderhaeghen, J.-J., Cholecystokinin receptors of A type in the human dorsal medulla oblongata and meningiomas, and of B type in small cell lung carcinomas, 243
Manfredi, M., see Cruccu, G., 68
Martellotta, M.C., see Demontis, M.G., 224
McKenzie, J.C., Cowie, R.J. and Inagami, T., ANP-like immunoreactivity in neuronal perikarya and processes associated with vessels of the pia and cerebral parenchyma in dog, 253
Melis, F., see Carboni, S., 129
Mesulam, M., see Schätz, C.R., 56
Miller, D.B., see O'Callaghan, J.P., 228
Moine, C.L., Tison, F. and Bloch, B., D_2 dopamine receptor gene expression by cholinergic neurons in the rat striatum, 248
Moore, M.A., see Strong, R., 175
Mori, Kenjiro, see Arai, T., 353
Mortilla, M., see Sorbi, S., 165
Morton, C.R. and Hutchison, W.D., Morphine does not reduce the intraspinal release of calcitonin gene-related peptide in the cat, 319
Mulholland, F., see Lasher, R.S., 43
Müller, F., see Stelmach, G.E., 188
Murakawa, M., see Arai, T., 353

Nabeshima, T., Tohyama, K. and Kameyama, T., Involvement of AChergic systems in chlordiazepoxide induced impairment of passive avoidance response, 200
Nakamura, Ya., see Kudo, M., 26
Nakao, S., see Arai, T., 353
Ng, J.P., see Brock, J.W., 234
Nitsch, R., see Hoyer, S., 358
O'Callaghan, J.P., Miller, D.B. and Reinhard, J.F.Jr., 1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)- induced damage of striatal dopaminergic fibers attenuates subsequent astrocyte response to MPTP, 228

Oesterreich, K., see Hoyer, S., 358
Ono, T., Fujino, Y., Tsuchiya, T. and Tsuda, M., Plasmid DNAs directly injected into mouse brain with lipofectin can be incorporated and expressed by brain cells, 259
Otten, U., see Gadient, R.A., 335

Pani, L., see Carboni, S., 129
Pardini, B.J., Lund, D.D. and Schmid, P.G., Innervation patterns of the middle cervical – stellate ganglion complex in the rat, 300
Phelix, C.F. and Hartle, D.K., Systemic glutamate induces degeneration of a subpopulation of serotonin-immunoreactive neurons in the area postrema of rats, 31
Piacentini, S., see Sorbi, S., 165
Piot, O., Betschart, J., Stutzmann, J.M. and Blanchard, J.C., Cyclopyrrolones, unlike some benzodiazepines, do not induce physical dependence in mice, 140
Populin, L., see Stelmach, G.E., 188
Puig, M.M., see Bansinath, M., 212

Rada, P. and Hernandez, L., Opposite changes of dopamine turnover in prefrontal cortex and nucleus accumbens after amygdaloid kindling, 144
Rahmann, H., see Evans, D.B.M.L., 117
Rajput, Ali, see Kish, S.J., 347
Ramabadran, K., see Bansinath, M., 212
Randić, M., Hećimović, H. and Ryu, P.D., Substance P modulates glutamate-induced currents in acutely isolated rat spinal dorsal horn neurones, 74
Rao, Z.-R., see Li, Yu.-Q., 285
Redle, S., see Hösli, L., 307

Reinhard, J.F.Jr., see O'Callaghan, J.P., 228
Réthelyi, M. and Lozsádi, D., Synapses upon the axon origin of dorsal horn neurons in the rat spinal cord, 20
Richardson, A., see Strong, R., 175
Rieger, M.N. and F., see Verdière-Sahuque, M., 341
Robitaille, Y., see Kish, S.J., 347
Rogers, A.T., Harrison, R., Lunt, G.G., Case, C.P. and Barraclough, C., Neuronotrophic activities of normal and motor neurone disease patient skeletal muscle extract on cultured rat spinal neurones, 194
Rojas, J., see Hösli, L., 307
Rossetti, Z.L., see Carboni, S., 129
Ryu, P.D., see Randić, M., 74

Saito, H., see Abe, K., 87
Saitoh, T., see Kametani, F., 62
Sampson, S.R., see Brodie, C., 325
Sanderson, R., see Lasher, R.S., 43
Sarda, N., see Houdouin, F., 218
Satoh, M., see Kaneko, S., 134
Sato, M., see Kametani, F., 62
Schätz, C.R., Geula, C. and Mesulam, M., Competitive substrate inhibition in the histochemistry of cholinesterase activity in Alzheimer's disease, 56
Scherman, D., see Leroux-Nicollet, I., 1
Schmid, P.G., see Pardini, B.J., 300
Schramek, H., see Hösli, L., 307
Shibata, R., see Kurata, K., 264
Shi, J.-W., see Li, Y.-Q., 285
Shinoda, T., see Kametani, F., 62
Shinoda, Y., see Kakei, S., 280
Shupliakov, O.V., Antal, M. and Székely, G., Recurrent dorsal root potentials and motoneuron morphology in the frog spinal cord, 289
Skaper, S.D., Facci, L. and Leon, A., Gangliosides attenuate the delayed neurotoxicity of aspartic acid in vitro, 154
Smith, P.A., see Zidichouski, J.A., 123
So, K.-F., see Tan, M.M.L., 14
Sontag, K. -H., see Jaspers, R.M.A., 149
Sorbi, S., Mortilla, M., Piacentini, S., Tonini, S. and Amaducci, L., Altered hexokinase activity in skin cultured fibroblasts and leukocytes from Alzheimer's disease patients, 165
Spuler, A., see Knöpfel, T., 295
Stelmach, G.E., Populin, L. and Müller, F., Postural muscle onset and voluntary movement in the elderly, 188
Stewart, John M., see Lasher, R.S., 43
Strong, R., Hale, C., Moore, M.A., Wessels-Reiker, M., Armbrecht, H.J. and Richardson, A., GABA receptor modulation of tyrosine hydroxylase gene expression in the rat adrenal gland, 175
Stutzmann, J.M., see Piot, O., 140
Sugimura, M., see Kaneko, S., 134
Székely, G., see Shupliakov, O.V., 289

Takahashi, Y., see Goto, K., 269
Tanaka, K., see Kametani, F., 62
Tan, M.M.L., Harvey, A.R. and So, K.-F., Regeneration of retinal axons in grafts of peripheral and central nervous tissue in the adult rat, 14
Tison, F., see Moine, C.L., 248
Tohyama, K., see Nabeshima, T., 200
Tonini, S., see Sorbi, S., 165
Tsuchiya, T., see Ono, T., 259
Tsuda, M., see Ono, T., 259
Turndorf, H., see Bansinath, M., 212

Urschel, B.A. and Hulsebosch, C.E., In vivo recovery of Schwann cell–neuronal interactions altered by nerve growth factor suppression in the rat, 50

Vanderhaeghen, J.-J., see Mailleux, P., 243
Verdière-Sahuque, M., Goudou, D., Garcia, L. and Rieger, M.N. and F., Cell modulation of hydrophobic tailed 16S acetylcholinesterase by intracellular calcium in rat superior cervical ganglion neurons, 341
Vulfius, C.A., see Erdélyi, L., 99

Walz, W., see Wuttke, W.A., 105
Wan, W. and Geiger, J.D., Effects of enucleations and visual deprivation on adenosine A_1 and A_2 receptor binding in superior colliculus of rat, 160
Watanabe, Ya., see Abe, K., 87
Wessels-Reiker, M., see Strong, R., 175
White, F.J., see Ackerman, J.M., 181
Wolf, G., Keilhoff, G., Fischer, S. and Hass, P., Subcutaneously applied magnesium protects reliably against quinolinate-induced *N*-methyl-D-aspartate (NMDA)-mediated neurodegeneration and convulsions in rats: are there therapeutical implications?, 207
Wuttke, W.A. and Walz, W., Sodium- and bicarbonate-independent regulation of intracellular pH in cultured mouse astrocytes, 105

Xie, F., see Abe, K., 87

Yamamoto, M. and Kondo, H., Gene expression of a neuronal growth-associated protein, GAP-43, in the paraganglionic carotid body as well as in the autonomic ganglia of normal adult rats, 275

Yu, S.P. and Der Kloot, W.V., Non-quantal acetylcholine release at mouse neuromuscular junction: effects of elevated quantal release and aconitine, 111

Zidichouski, J.A., Chen, H. and Smith, P.A., Neuropeptide Y activates inwardly-rectifying K-channels in C-cells of amphibian sympathetic ganglia, 123